Atomic Theory Review

Fill in t	he Blank						
	1. The number of pro	The number of protons and neutrons in the nucleus of an atom is the					
	2. Atoms possessing t	. Atoms possessing the same number of protons but a different number of neutrons are					
	3. The lowest energy						
	4. A particle found in						
	5. The number of pro	 5. The number of protons in an atom is represented by the 6. The identity of an element is determined by the number of present. 7. The volume and shape of the region in space where you would be most likely to locate an electron is a(n) 					
	6. The identity of an						
	•						
	8. A system for predi	8. A system for predicting the order of filling energy sublevels with electrons is the					
2.	Of the following assumptions or reatomic theory, the only one that recorrect in most cases is A) All atoms of an element are ident B) Atoms are indivisible and indestra C) Oxygen has an atomic weight of 7 D) Atoms of elements combine in the whole numbers to form compound Rutherford's experiments on the particles by thin metal foils established A) The mass and charge of an atom in a nucleus. B) Electrons are fundamental particles C) All electrons have the same charge C) Atoms are electrically neutral.	remains essentially rical in mass. Luctible.	$^{32}_{16}S$ is A) $^{35}_{17}Cl^{-}$ B) $^{34}_{16}S^{+}$ C) $^{40}_{18}Ar^{2+}$ D) $^{35}_{16}S^{2-}$ All of the following masses are possible for an individual carbon atom except one. That impossible one is A) 12.000 00 amu C) 12.011 15 amu B) 13.003 35 amu D) 14.003 24 amu				

Short Answer

- Compare the similarities and differences between Rutherford's and Bohr's theories of the atom.
- Compare the similarities and differences between Bohr's model and the Quantum Mechanical model of the atom.
- 3. Draw energy level diagrams, and write electron configurations and energy level populations for lead and Rutherfordium.
- 4. Complete the following table.

Symbol	Atomic Number	Mass Number	Protons	Neutrons	Electrons
38 18 A r					
¹⁸ ₈ O					
³⁶ ₁₆ S ²⁻					
X-				20	18
		26		14	10

5. Calculate the average atomic mass of nickel that has five naturally occurring isotopes: 67.88% nickel-58 (mass = 57.9353 u), 26.23% nickel 60 (mass = 58.9332 u), 1.19%

- nickel-61 (mass = 60.9310 u), 3.66% nickel-62 (mass = 61.9283 u), and 1.08% nickel-64 (mass = 63.9280 u).
- 6. There are two principal isotopes of silver, $^{107}_{47}$ Ag and $^{109}_{47}$ Ag. The atomic weight of naturally occurring silver is 107.87, with 51.82% of the atoms being $^{107}_{47}$ Ag. The mass of an atom of $^{107}_{47}$ Ag is 106.9 amu. What is the mass, in amu, of an atom of $^{109}_{47}$ Ag?
- Naturally occurring indium consists of two isotopes: indium-113 with an atomic mass of 112.9043 u, and indium-115 with an atomic mass of 114.9041 u. If the average atomic mass of indium is 114.82 u, calculate the percentage of each isotope of naturally occurring indium.
- *8. Naturally, occurring zinc consists of five isotopes, ⁶⁴Zn, ⁶⁶Zn, ⁶⁷Zn, ⁶⁸Zn, and ⁷⁰Zn, whose atomic masses are 63.929, 65.926, 66.927, 67.925, and 69.925 amu, respectively. The most abundant isotopes are ⁶⁴Zn, ⁶⁶Zn, and ⁶⁸Zn, which accounts for 48.89%, 27.81%, and 18.57%, respectively, of naturally occurring zinc. Given that the observed atomic mass of zinc is 65.38 amu, calculate the percentages of the two remaining isotopes of zinc.
- *9. Give all possible sets of quantum numbers for an electron in a 4f orbital.